



Cardiac Myocyte Medium-serum free (CMM-sf) Catalog #6101

Product Description

Cardiac Myocyte Medium-serum free (CMM-sf), when used with Cardiac Myocyte Growth Supplement-serum free (CMGS-sf, Cat #6152) is a complete medium designed for the culture of normal cardiac myocytes in serum-free conditions. It is a sterile, liquid medium which contains essential and non-essential amino acids, vitamins, organic and inorganic compounds, hormones, growth factors and trace minerals. The medium is HEPES and bicarbonate buffered and has a pH of 7.4 when equilibrated in an incubator with an atmosphere of 5% CO₂/95% air. The medium is formulated (quantitatively and qualitatively) to provide an optimally balanced nutritional environment that selectively supports the culture of normal cardiac myocytes *in vitro* under serum free conditions.

Components

CMM-sf consists of 500 ml of basal medium, 5 ml of Cardiac Myocyte Growth Supplement-serum free (CMGS-sf, Cat. #6152) and 5 ml of penicillin/streptomycin solution (P/S, Cat. #0503). *Note: CMGS-sf and P/S are not pre-mixed in CMM-sf; they must be added separately to make the complete CMM-sf.*

Product Use

CMM-sf is for research use only. It is not approved for human or animal use, or for application in *in vitro* diagnostic procedures.

Storage

Store the basal medium at 4°C and the CMGS-sf and P/S solution at -20°C. Protect from light.

Shipping

Basal medium: room temperature. Supplements: dry ice.

Instructions for Use

Thaw CMGS-sf and P/S solution at 37°C. Gently tilt the tubes several times to ensure the contents are completely mixed before adding to the medium. Spray the medium bottle and tubes with 70% ethanol, and wipe to remove excess liquid. In a sterile field, remove the caps without touching the interior threads with fingers. Add CMGS-sf and P/S solution to the medium and mix well. Since several components are light-labile, the medium should not be exposed to light for extended periods. We do not recommend warming medium in a 37°C water bath prior to use. When stored in the dark at 4°C, the reconstituted medium is stable for one month.

Caution: If handled improperly, some components of the medium may present a health hazard. Take appropriate precautions when handling it, including the wearing of protective clothing and eyewear. Dispose of properly.